

FERMAID O™

Organic yeast nutrient

DESCRIPTION

FERMAID O™ is a nutrient resulting from Lallemand's research on nitrogen metabolism during fermentation.

FERMAID O™ is a unique yeast autolysate with a high content of free amino-acids and peptides, selected by Lallemand. It contains 100% of organic nitrogen and therefore does not contain any ammonia salts (DAP or SDA).



BENEFITS & RESULTS

FERMAID O™ supplies well balanced nutrients for yeasts:

- highly available amino-acids and peptides. Amino-acids are more efficiently used by yeast than inorganic nitrogen.
- natural sources of survival factors to help yeast in stressed conditions.
- natural sources of micronutrients such as vitamins (thiamine, biotine, panthotenic acid...) and oligoelements (magnesium, manganese, zinc...).

FERMAID O™ enhances ripe fruit character and volume while decreasing dryness sensation.

With FERMAID O™, nitrogen is assimilated more steadily than mineral nitrogen preventing fermentation activity peak while limiting temperature peaks. Note: in low nutrient situations, yeast assimilable nitrogen (YAN) may be sufficient to avoid fermentation problems.




Figure 1: Addition of 16 mg/L of YAN at 2 stages of fermentation (at the beginning and at 1/3 of the alcoholic fermentation -AF-) in 2 different forms: inorganic nitrogen (DAP) and organic nitrogen (FERMAID O™).

On the figure 1, it is shown that for an equivalent amount of assimilable nitrogen added, the addition of organic nitrogen is efficient on the fermentative kinetics. Indeed, in a must with high nitrogen deficiency, it allows to consume all sugars whereas the same amount of YAN provided in the form of inorganic nitrogen does not allow to complete the fermentation. No difference can be noted between the control fermentation (no nutrient added) and the fermentation with added DAP addition. In those two cases, a stuck fermentation is observed.




Figure 2: Effect of inorganic/organic nutrition on a Chardonnay fermented at 16°C (AWRI, 2009).
DAP at 50 mg/L YAN on aromas compounds production.




Figure 3: Effect of FERMAID O™ on the sensory profile of a Syrah (ICV R&D Dpt, 2007).

INSTRUCTIONS FOR OENOLOGICAL USE

	YAN (Yeast Assimilable Nitrogen) in mg/L	
	30 g/hL added product	40 g/hL added product
FERMAID O™	36 mg/L	48 mg/L
DAP	63 mg/L	84 mg/L

Recommended dosage: 2x20g/hL (to supply the must with 15 - 20mg/L organic YAN)

- 20 g/hL at the beginning of AF
- 20 g/hL at 1/4 to 1/3 of AF
- Suspend in 10 times its weight of water or must and add to the must during alcoholic fermentation.



OMRI (Organic Materials Review Institute) is a US national nonprofit organization that determines which input products are allowed for use in organic production and processing.

PACKAGING AND STORAGE

- 10 kg (4 x 2.5 kg bags) and 10 kg box.
- Store in a cool dry place.
- To be used once opened.

Distributed by:

The information in this document is correct to the best of our knowledge. However, this data sheet should not be considered to be an express guarantee, nor does it have implications as to the sales condition of this product. June 2022.



LALLEMAND OENOLOGY
Original by culture

www.lallemantwine.com

Visionary biological solutions - Being original is key to your success. At Lallemand Oenology, we apply our passion for innovation, maximize our skill in production and share our expertise, to select and develop natural microbiological solutions. Dedicated to the individuality of your wine, we support your originality, we cultivate our own.