

LALVIN® EC 1118®

ORIGIN AND APPLICATION

Robust, Reliable and Neutral. Useful for a wide range of applications, including wine and fruit cider fermentations.


A yeast selected in the Champagne region for its excellent properties in producing base wine for Champagne as well as “in-bottle” secondary fermentation.

Lalvin EC 1118® is known for its robust and reliable fermentation kinetics. Its sensory contribution is considered neutral, as it gives very little yeast sensory contribution to the wine.

It is used extensively in the world for the production both white and red wines.



MICROBIAL AND OENOLOGICAL PROPERTIES

- Recommended for white, rose and red wine production. Highly recommended for secondary fermentation. 
- *Saccharomyces cerevisiae* var. *bayanus*
- Desirable fermentation temperature: 10-30°C.
- Alcohol tolerance 18% v/v *subject to fermentation conditions.
- Low relative nitrogen demand (under controlled laboratory conditions)
- Short lag phase and high fermentation vigour. Cooling may be required to control this high vigour.
- Very low production of H₂S under low YAN conditions
- Low production of SO₂ binding compounds.
- Moderate relative potential for SO₂ production (can produce high levels of SO₂ under low nutrient conditions, up to 50mg/L). Generally considered to be neutral to MLF.
- Killer factor active.
- Low foam producer.

INSTRUCTION FOR USE

Dosage Rate:

- 25g/hL of Active Dried Yeast (this will provide an initial cell population of approximately 5 x10⁶ viable cells/ml)
- 30g/hL of Go-Ferm Protect Evolution™
- Nitrogen source from the Fermaid™ range

INSTRUCTION FOR USE (cont'd)

Procedure for 1000L ferment.

- 1) Add 300g of Go-Ferm Protect Evolution™ to 5L of 40-43°C clean, chlorine free water. Stir until an homogenous suspension free of lumps is achieved.
- 2) When the temperature of this suspension is between 35-40°C, sprinkle 250g of yeast slowly and evenly onto the surface of the water, whilst gently stirring. Ensure any clumps are dispersed.
- 3) Allow to stand for 20 minutes before further gently mixing.
- 4) Mix the rehydrated yeast with a little juice, gradually adjusting the yeast suspension temperature to within 5-10°C of the juice/must temperature.
- 5) Inoculate into the must.

Further Notes

- Steps 1-5 should be completed within 30 minutes.
- It is best to limit first juice/must volume addition to one tenth the yeast suspension volume and wait 10 minutes before the addition to juice.
- To minimize cold shock, ensure temperature changes are less than 10°C.
- It is recommended that juice / must be inoculated no lower than 18°C.
- It is recommended to use complex nutrition nitrogen source, such as either **Fermaid K™** or **Fermaid O™**.

PACKAGING AND STORAGE

- All Active Dried Yeast should be stored dry, between 4-12°C and the vacuum packaging should remain intact.

The information herein is true and accurate to the best of our knowledge; however, this data sheet is not to be considered as a guarantee, expressed or implied, or as a condition of sale of this product.



WINE
YEASTS



WINE
BACTERIA



NUTRIENTS
/PROTECTORS



SPECIFIC
INACTIVATED YEASTS



ENZYMES



CHITOSAN



VINEYARD
SOLUTIONS

LALLEMAND

LALLEMAND OENOLOGY

Original *by culture*